




# Cosmology with the SKA

Chris Blake (Swinburne)

# Our current model of cosmology

- We have a superbly detailed picture of the **early Universe** [e.g. CMB, nucleosynthesis]
- We have a model for the **evolution of the Universe** that matches a range of cosmological data
- This model invokes 3 new pieces of physics : **inflation**, **dark matter** and **dark energy**





# Dark energy : is it vacuum energy?


A cosmological constant matches the data so far, but its amplitude is inexplicable



It's just SOMETHING  
You'll never  
understand...



© Bill Junk 2008




# Cosmology : the optimistic viewpoint !

- Dark matter and energy show that our understanding of physics is incomplete
- Astronomy can provide fundamental physical insights into quantum theory, gravity, and particle physics
- We are working in the breakthrough era where new data should be revolutionary!



# Large-scale structure

- **Geometrical information**  
(e.g. baryon acoustic oscillations)
- **Gravitational information**  
(e.g. redshift-space distortions)
- **Primordial information**  
(e.g. shape of large-scale power spectrum)





# Cosmology with the SKA

- 2004 style ...

ζiv:astro-ph/0409278v1 13 Sep 2004

## Cosmology with the SKA

C.A. Blake<sup>a</sup>, F.B. Abdalla<sup>b</sup>, S.L. Bridle<sup>c</sup>, S. Rawlings<sup>b</sup>

<sup>a</sup>School of Physics, University of New South Wales, Sydney, NSW 2052, Australia

<sup>b</sup>Astrophysics, Department of Physics, University of Oxford, Keble Road, Oxford, OX1 3RH, UK

<sup>c</sup>Department of Physics and Astronomy, University College London, London, WC1E 6BT, UK

We argue that the Square Kilometre Array has the potential to make both redshift (HI) surveys and radio continuum surveys that will revolutionize cosmological studies, provided that it has sufficient instantaneous field-of-view that these surveys can cover a hemisphere ( $f_{\text{sky}} \sim 0.5$ ) in a timescale  $\sim 1$  yr. Adopting this assumption, we focus on two key experiments which will yield fundamental new measurements in cosmology, characterizing the properties of the mysterious dark energy which dominates the dynamics of today's Universe. Experiment I will map out  $\sim 10^9 (f_{\text{sky}}/0.5)$  HI galaxies to redshift  $z \approx 1.5$ , providing the premier measurement of the clustering power spectrum of galaxies: accurately delineating the acoustic oscillations and the 'turnover'. Experiment II will quantify the cosmic shear distortion of  $\sim 10^{10} (f_{\text{sky}}/0.5)$  radio continuum sources, determining a precise power spectrum of the dark matter, and its growth as a function of cosmic epoch. We contrast the performance of the SKA in precision cosmology with that of other facilities which will, probably or possibly, be available on a similar timescale. We conclude that data from the SKA will yield transformational science as the direct result of four key features: (i) the immense cosmic volumes probed, exceeding future optical redshift surveys by more than an order of magnitude; (ii) well-controlled systematic effects such as the narrow ' $k$ -space window function' for Experiment I and the accurately-known 'point-spread function' (synthesized beam) for Experiment II; (iii) the ability to measure with high precision large-scale modes in the clustering power spectra, for which nuisance effects such as non-linear structure growth, peculiar velocities and 'galaxy bias' are minimised; and (iv) different degeneracies between key parameters to those which are inherent in the Cosmic Microwave Background.

# Cosmology with the SKA

- 2015 style ...

## Cosmology with the SKA – overview

---

**Roy Maartens<sup>\*1,2</sup>, Filipe B. Abdalla<sup>3,4</sup>, Matt Jarvis<sup>5,1</sup>, Mario G. Santos<sup>1,6</sup> –  
on behalf of the SKA Cosmology SWG**

<sup>1</sup> *Department of Physics, University of Western Cape, Cape Town 7535, South Africa*

<sup>2</sup> *Institute of Cosmology & Gravitation, University of Portsmouth, Portsmouth PO1 3FX, UK*

<sup>3</sup> *Department of Physics & Astronomy, University College London, London WC1E 6BT, UK*

<sup>4</sup> *Department of Physics & Electronics, Rhodes University, Grahamstown 6140, South Africa*

<sup>5</sup> *Astrophysics, University of Oxford, Oxford OX1 3RH, UK*

<sup>6</sup> *SKA South Africa, The Park, Park Road, Cape Town 7405, South Africa*

*E-mail: [roy.maartens@gmail.com](mailto:roy.maartens@gmail.com)*





The new frontier of cosmology will be led by three-dimensional surveys of the large-scale structure of the Universe. Based on its all-sky surveys and redshift depth, the SKA is destined to revolutionize cosmology, in combination with future optical/ infrared surveys such as Euclid and LSST. Furthermore, we will not have to wait for the full deployment of the SKA in order to see transformational science. In the first phase of deployment (SKA1), all-sky HI intensity mapping surveys and all-sky continuum surveys are forecast to be at the forefront on the major questions of cosmology. We give a broad overview of the major contributions predicted for the SKA. The SKA will not only deliver precision cosmology – it will also probe the foundations of the standard model and open the door to new discoveries on large-scale features of the Universe.

# Cosmology with the SKA

- Use **HI intensity mapping** to measure large-scale structure across large volumes to high redshift
- Test cosmology with **cross-correlations** of the radio continuum survey with CMB and optical surveys
- Measure **weak gravitational lensing** in radio continuum observations using new techniques
- Constrain **primordial non-Gaussianity** using the largest-scale modes




# Upcoming optical surveys



Euclid


# HI intensity mapping

- 21cm surveys which do not detect individual galaxies but the integrated emission in each pixel of a datacube




Chang et al. (2008)

Masui et al. (2013)




# HI intensity mapping


- Great potential owing to the volume probed!



## Dark energy measurements:



## BAO distance/expansion measurements:





Bull et al. (2014)



# HI intensity mapping

- Challenges : enormous foreground subtraction problem, including polarization leakage




- Need realistic simulations!

See Laura's talk ...

# Continuum surveys : cross-correlations



- Radio continuum surveys trace a high- $z$  density map
- Expect correlations with CMB (late-time ISW effect and lensing) and low- $z$  galaxies (magnification)



Planck collaboration  
(2013)

# Continuum surveys : late-time ISW effect

- Cross-correlation between the NVSS and CMB



Catalogue	$A \pm \sigma_A$	$S/N$	expected $S/N$
2MASS cut	$1.40 \pm 2.09$	0.7	0.5
SDSS gal DR8	$1.24 \pm 0.57$	2.2	1.6
SDSS LRG DR7	$2.10 \pm 0.84$	2.5	1.2
NVSS	$1.21 \pm 0.43$	2.8	2.6
HEAO	$1.37 \pm 0.56$	2.4	2.0
SDSS QSO DR6	$1.43 \pm 0.62$	2.3	1.7
<b>TOTAL</b>	<b><math>1.38 \pm 0.32</math></b>	<b><math>4.4 \pm 0.4</math></b>	<b><math>\approx 3.1, &lt; 7.6</math></b>

Nolta et al. (2004)



Ho et al. (2008)

Giannantonio et al. (2013)




# Continuum surveys : late-time ISW effect

- Physical evidence for dark energy






Granett et al. (2008)  
de Putter et al. (2010)

- Problem : limited precision





# Continuum surveys : gravitational lensing

- Probes total mass distribution and spacetime metric




# Continuum surveys : gravitational lensing

- Detection of cosmic shear in FIRST survey



Chang et al. (2004)






# Continuum surveys : gravitational lensing


- What is the potential radio advantage?

Probing to **high redshifts**




**PSF calibration** a challenge in optical


Shape estimates with **uncorrelated systematics** between radio/optical



**Estimates of intrinsic shape** using polarization and/or velocity maps




Morales (2006)



Brown & Battye (2010)

# Continuum surveys : gravitational lensing



- What are the challenges?



Credit : <http://radiogreat.jb.man.ac.uk>



# Continuum surveys : non-Gaussianity

- Largest-scale structure contains primordial imprint



Ferramacho  
et al. (2014)



Seljak (2009)





# Continuum surveys : homogeneity and isotropy



- Are CMB “anomalies” cosmologically interesting?



Test with **radio**  
**galaxy dipole**

Blake & Wall (2002)

Rubart & Schwarz (2013)



# Cosmology with the SKA

- Use **HI intensity mapping** to measure large-scale structure across large volumes to high redshift
- Test cosmology with **cross-correlations** of the radio continuum survey with CMB and optical surveys
- Measure **weak gravitational lensing** in radio continuum observations using new techniques
- Constrain **primordial non-Gaussianity** using the largest-scale modes